

Hybrid High-Order methods

Introduction and fully discrete analysis framework

Daniele A. Di Pietro



22 May 2024

Outline

- 1 Preliminaries
- 2 A non-conforming finite element scheme on standard meshes
- 3 An Hybrid High-Order scheme on polytopal meshes
- 4 Fully discrete analysis framework

Outline

- 1 Preliminaries
- 2 A non-conforming finite element scheme on standard meshes
- 3 An Hybrid High-Order scheme on polytopal meshes
- 4 Fully discrete analysis framework

Setting

- Let $\Omega \subset \mathbb{R}^d$, $d \geq 2$, be an open connected polytopal domain
- We focus on the Poisson problem: Given $f : \Omega \rightarrow \mathbb{R}$, find $u : \Omega \rightarrow \mathbb{R}$ s.t.

$$\begin{aligned}-\Delta u &= f && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega\end{aligned}$$

- Let $f \in L^2(\Omega)$. A possible weak formulation reads: Find $u \in H_0^1(\Omega)$ s.t.

$$a(u, v) := \int_{\Omega} \nabla u \cdot \nabla v = \int_{\Omega} f v \quad \forall v \in H_0^1(\Omega)$$

- The well-posedness of this problem hinges on the **Poincaré inequality**

$$\|v\|_{L^2(\Omega)} \leq C_{\Omega} \|\nabla v\|_{L^2(\Omega)^d}$$


Local polynomial spaces and L^2 -orthogonal projector

- Denote by \mathbb{P}_d^ℓ the space of d -variate polynomials of total degree $\leq \ell$
- Let $Y \subset \mathbb{R}^d$ and denote by $\mathcal{P}^\ell(Y)$ the restriction of \mathbb{P}_d^ℓ to Y
- Given $v \in L^2(Y)$, its L^2 -orthogonal projection on $\mathcal{P}^\ell(Y)$ is s.t.

$$\int_Y (v - \pi_{\mathcal{P}^\ell(Y)} v) w = 0 \quad \forall w \in \mathcal{P}^\ell(Y)$$

- $\pi_{\mathcal{P}^\ell(Y)} v$ optimally approximates v in all Sobolev seminorms under mild assumptions on Y ; see [DP and Droniou, 2020, Chapter 1] for details

Raviart–Thomas–Nédélec and Crouzeix–Raviart elements I




- Denote by T a d -simplex and by \mathcal{F}_T the set collecting its faces
- Let $\mathcal{RTN}^1(T) := \mathcal{P}^0(T)^d + x\mathcal{P}^0(T)$
- Define the degrees of freedom $\sigma := (\sigma_F)_{F \in \mathcal{F}_T}$ s.t.

$$\sigma_F : H^1(T)^d \ni \tau \mapsto \frac{1}{|F|} \int_F \tau \cdot n_{TF} \in \mathbb{R}$$

- Notice that $\tau \cdot n_{TF} \in \mathcal{P}^0(F)$ for all $\tau \in \mathcal{RTN}^l(T)$ and all $F \in \mathcal{F}_T$
- $(T, \mathcal{RTN}^1(T), \sigma)$ is a FE [Raviart and Thomas, 1977, Nédélec, 1980]

Raviart–Thomas–Nédélec and Crouzeix–Raviart elements II



- Let now $\mathcal{P}^1(T)$ be the space of affine functions on T
- Define the degrees of freedom $\sigma := (\sigma_F)_{F \in \mathcal{F}_T}$ s.t.


$$\sigma_F : H^1(T) \ni v \mapsto \frac{1}{|F|} \int_F v \in \mathbb{R}$$

- $(T, \mathcal{P}^1(T), \sigma)$ is a FE [Crouzeix and Raviart, 1973]

Outline

- 1 Preliminaries
- 2 A non-conforming finite element scheme on standard meshes
- 3 An Hybrid High-Order scheme on polytopal meshes
- 4 Fully discrete analysis framework

A non-conforming finite element scheme I



- Let \mathcal{T}_h be a conforming simplicial mesh of Ω
- Let $C\mathcal{R}(\mathcal{T}_h) \subset H^1(\mathcal{T}_h)$ be the Crouzeix–Raviart space on \mathcal{T}_h and set

$$C\mathcal{R}_0(\mathcal{T}_h) := \{v_h \in C\mathcal{R}(\mathcal{T}_h) : \pi_{\mathcal{P}^0(F)} v = 0 \text{ for all } F \in \mathcal{F}_h^b\}$$

- Notice that $C\mathcal{R}_0(\mathcal{T}_h) \not\subset H_0^1(\Omega)$!
- With ∇_h **broken gradient**, the scheme reads: Find $u_h \in C\mathcal{R}_0(\mathcal{T}_h)$ s.t.

$$a_h(u_h, v_h) := \int_{\Omega} \nabla_h u_h \cdot \nabla_h v_h = \int_{\Omega} f v_h \quad \forall v_h \in C\mathcal{R}_0(\mathcal{T}_h)$$

Stability analysis I

Lemma (Discrete Poincaré inequality in the Crouzeix–Raviart space)

For all $v_h \in \mathcal{CR}_0(\mathcal{T}_h)$,

$$\|v_h\|_{L^2(\Omega)} \lesssim \|\nabla_h v_h\|_{L^2(\Omega)^d}$$

where $a \lesssim b$ means $a \leq Cb$ with C independent of h .

- The Poincaré–Wirtinger inequality gives, setting $v_T := (v_h)|_T$,

$$\|v_T\|_{L^2(T)} \lesssim \|\pi_{\mathcal{P}^0(T)} v_T\|_{L^2(T)} + h_T \|\nabla v_T\|_{L^2(T)^d}$$

- Hence, letting $\bar{v}_h := \pi_{\mathcal{P}^0(\mathcal{T}_h)} v_h$, it suffices to prove that

$$\|\bar{v}_h\|_{L^2(\Omega)} \lesssim \|\nabla_h v_h\|_{L^2(\Omega)^d}$$

- Since $\operatorname{div} : \mathcal{RTN}^1(\mathcal{T}_h) \rightarrow \mathcal{P}^0(\mathcal{T}_h)$ is surjective: $\exists \tau_h \in \mathcal{RTN}^1(\mathcal{T}_h)$ s.t.

$$\operatorname{div} \tau_h = \bar{v}_h \text{ and } \|\tau_h\|_{H(\operatorname{div}; \Omega)} \lesssim \|\bar{v}_h\|_{L^2(\Omega)}$$

Stability analysis II

- We write, letting $v_T := (v_h)_{|T}$ for all $T \in \mathcal{T}_h$,

$$\|\bar{v}_h\|_{L^2(\Omega)}^2 \stackrel{\triangle}{=} \sum_{T \in \mathcal{T}_h} \int_T v_T \operatorname{div} \tau_h \stackrel{\text{IBP}}{=} - \int_{\Omega} \nabla_h v_h \cdot \tau_h + \sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} \int_F v_T (\tau_h \cdot n_{TF})$$

- Replacing $v_T \leftarrow \pi_{\mathcal{P}^0(F)} v_T =: \textcolor{red}{v}_F$ in the boundary term and rearranging,


$$\begin{aligned} \sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} \int_F v_T (\tau_h \cdot n_{TF}) &= \sum_{F \in \mathcal{F}_h} \sum_{T \in \mathcal{T}_F} \int_F v_T (\tau_h \cdot n_{TF}) \\ &= \sum_{F \in \mathcal{F}_h^i} \sum_{T \in \mathcal{T}_F} \int_F \textcolor{red}{v}_F (\tau_h \cdot n_{TF}) + \sum_{F \in \mathcal{F}_h^b} \int_F \textcolor{red}{v}_F (\tau_h \cdot n_F) = 0, \end{aligned}$$

by single-valuedness of v_F for $F \in \mathcal{F}_h^i$ and $v_F = 0$ for $F \in \mathcal{F}_h^b$

- Using Cauchy–Schwarz inequalities, we get

$$\|\bar{v}_h\|_{L^2(\Omega)}^2 \leq \|\nabla_h v_h\|_{L^2(\Omega)^d} \|\tau_h\|_{L^2(\Omega)^d} \lesssim \|\nabla_h v_h\|_{L^2(\Omega)^d} \|\bar{v}_h\|_{L^2(\Omega)}$$

Limitations of the finite element approach



- Approach limited to **conforming** meshes with **standard elements**
 - ⇒ local refinement requires to **trade mesh size for mesh quality**
 - ⇒ complex geometries may require a **large number of elements**
 - ⇒ the element shape cannot be **adapted to the solution**
- Treating more general meshes in the FE spirit would significantly increase the space dimension [Droniou et al., 2021]
- The extension to **high-order** is not straightforward

Fully discrete polytopal approach



- Key idea: replace both spaces and operators by discrete counterparts
- Support of **polyhedral meshes** and **high-order**
- Several strategies to **reduce the number of unknowns** on general shapes
- Elegant analysis framework available

A few key references

- Introduction of Hybrid High-Order (HHO) methods [DP et al., 2014]
- Fully discrete analysis framework [DP and Droniou, 2018]
- A monograph on HHO methods [DP and Droniou, 2020]
- Introduction of Discrete de Rham (DDR) methods [DP et al., 2020]
- DDR for the de Rham complex of differential forms [Bonaldi et al., 2023]

Outline

- 1 Preliminaries
- 2 A non-conforming finite element scheme on standard meshes
- 3 An Hybrid High-Order scheme on polytopal meshes
- 4 Fully discrete analysis framework

A paradigm shift

- Let $v_T \in \mathcal{P}^1(T)$ and set $v_F := \pi_{\mathcal{P}^0(F)}(v_T)_{|F}$ for all $F \in \mathcal{F}_T$
- We have, for all $\tau \in \mathcal{P}^0(T)^d$,


$$\int_T \nabla v_T \cdot \tau = - \int_T v_T \operatorname{div} \tau + \sum_{F \in \mathcal{F}_T} \int_F v_T (\tau \cdot n_{TF}) = \sum_{F \in \mathcal{F}_T} \int_F v_F (\tau \cdot n_{TF})$$

- Moreover, with \bar{x}_Y center of mass of $Y \in \{T\} \cup \mathcal{F}_T$, noticing that $\bar{x}_T = \frac{1}{\operatorname{card}(\mathcal{F}_T)} \sum_{F \in \mathcal{F}_T} \bar{x}_F$ and using linearity,

$$\pi_{\mathcal{P}^0(T)} v = v_T(\bar{x}_T) = \frac{1}{\operatorname{card}(\mathcal{F}_T)} \sum_{F \in \mathcal{F}_T} v_T(\bar{x}_F) = \frac{1}{\operatorname{card}(\mathcal{F}_T)} \sum_{F \in \mathcal{F}_T} \frac{1}{|F|} \int_F v_F$$

- These formulas remain valid when T is a general polytope!

Generalization to polytopes




- Let $\underline{V}_T^0 := \{\underline{v}_T := (v_F)_{F \in \mathcal{F}_T} : v_F \in \mathcal{P}^0(F) \text{ for all } F \in \mathcal{F}_T\}$
- We can define a **potential reconstruction** $r_T^1 : \underline{V}_T^0 \rightarrow \mathcal{P}^1(T)$ enforcing

$$\int_T \nabla r_T^1 \underline{v}_T \cdot \tau = \sum_{F \in \mathcal{F}_T} \int_F v_F(\tau \cdot n_{TF}) \quad \forall \tau \in \mathcal{P}^0(T)^d$$

$$\pi_{\mathcal{P}^0(T)}(r_T^1 \underline{v}_T) = \frac{1}{\text{card}(\mathcal{F}_T)} \sum_{F \in \mathcal{F}_T} \frac{1}{|F|} \int_F v_F$$

Extension to arbitrary-order I



- Let $k \geq 0$ and define the **Hybrid High-Order (HHO) space**


$$\underline{V}_T^k := \left\{ \underline{v}_T = (v_T, (v_F)_{F \in \mathcal{F}_T}) : v_T \in \mathcal{P}^{k-1}(T) \text{ and } v_F \in \mathcal{P}^k(F) \text{ for all } F \in \mathcal{F}_T \right\}$$

- We define $r_T^{k+1} : \underline{V}_T^k \rightarrow \mathcal{P}^{k+1}(T)^d$ s.t., for all $\underline{v}_T \in \underline{V}_T^k$,

$$\int_T \nabla r_T^{k+1} \underline{v}_T \cdot \nabla w = - \int_T v_T \Delta w + \sum_{F \in \mathcal{F}_T} \int_F v_F (\nabla w \cdot n_{TF}) \quad \forall w \in \mathcal{P}^{k+1}(T),$$
$$\pi_{\mathcal{P}^0(T)}(r_T^1 \underline{v}_T) = \pi_{\mathcal{P}^0(T)} v_T,$$

where $v_T := \frac{1}{\text{card}(\mathcal{F}_T)} \sum_{F \in \mathcal{F}_T} \frac{1}{|F|} \int_F v_F$ if $k = 0$

Global HHO space and H^1 -like seminorm



- Given a polytopal mesh \mathcal{T}_h of Ω , define the global HHO space

$$\underline{V}_h^k := \left\{ \underline{v}_h = ((v_T)_{T \in \mathcal{T}_h}, (v_F)_{F \in \mathcal{F}_T}) \in \underline{V}_h^k : \right. \\ \left. v_T \in \mathcal{P}^{k-1}(T) \text{ for all } T \in \mathcal{T}_h \text{ and } v_F \in \mathcal{P}^k(F) \text{ for all } F \in \mathcal{F}_h \right\}$$

- We define on \underline{V}_h^k the norm

$$\|\underline{v}_h\|_{1,h}^2 := \sum_{T \in \mathcal{T}_h} \|\underline{v}_T\|_{1,T}^2$$

where $\|\underline{v}_T\|_{1,T}^2 := \|\nabla v_T\|_{L^2(T)^d}^2 + h_T^{-1} \sum_{F \in \mathcal{F}_T} \|v_F - v_T\|_{L^2(F)}^2$ for all $T \in \mathcal{T}_h$

Discrete Poincaré inequality in HHO spaces I

Lemma (Poincaré inequality in HHO spaces)

Denote by $\underline{V}_{h,0}^k$ the subspace of \underline{V}_h^k with vanishing boundary values. For any $\underline{v}_h \in \underline{V}_{h,0}^k$, letting $v_h \in \mathcal{P}^{\max(k-1,0)}(\mathcal{T}_h)$ be s.t. $(v_h)|_T := v_T$ for all $T \in \mathcal{T}_h$, it holds

$$\|v_h\|_{L^2(\Omega)} \lesssim \|\underline{v}_h\|_{1,h},$$

hence $\|\cdot\|_{1,h}$ is a norm on $\underline{V}_{h,0}^k$.

- Since $\operatorname{div} : H^1(\Omega)^d \rightarrow L^2(\Omega)$ is surjective, there is $\tau \in H^1(\Omega)^d$ s.t.

$$\operatorname{div} \tau = v_h \text{ and } \|\tau\|_{H^1(\Omega)^d} \lesssim \|v_h\|_{L^2(\Omega)}$$

- Notice that here we cannot seek τ in $\mathcal{RTN}^{k+1}(\mathcal{T}_h)$ since \mathcal{T}_h is not conforming simplicial!

Discrete Poincaré inequality in HHO spaces II

- We go on writing

$$\begin{aligned} \|v_h\|_{L^2(\Omega)}^2 &= \int_{\Omega} v_h \operatorname{div} \tau = \sum_{T \in \mathcal{T}_h} \int_T v_T \operatorname{div} \tau \\ &\stackrel{\text{IBP}}{=} \sum_{T \in \mathcal{T}_h} \left(- \int_T \nabla v_T \cdot \tau + \sum_{F \in \mathcal{F}_T} \int_F (v_T - \underline{v}_F)(\tau \cdot n_{TF}) \right) \\ &\leq \left[\sum_{T \in \mathcal{T}_h} \left(\|\nabla v_T\|_{L^2(T)^d}^2 + h_T^{-1} \sum_{F \in \mathcal{F}_T} \|v_T - v_F\|_{L^2(F)}^2 \right) \right]^{\frac{1}{2}} \\ &\quad \times \left[\sum_{T \in \mathcal{T}_h} \left(\|\tau\|_{L^2(T)^2}^2 + h_T \|\tau\|_{L^2(\partial T)}^2 \right) \right]^{\frac{1}{2}} \end{aligned}$$

- Recalling the definition of $\|\cdot\|_{1,h}$ and using trace inequalities along with $h_T \leq h_\Omega \lesssim 1$, we get

$$\|v_h\|_{L^2(\Omega)}^2 \lesssim \|\underline{v}_h\|_{1,h} \|\tau\|_{H^1(\Omega)^d} \lesssim \|\underline{v}_h\|_{1,h} \|v_h\|_{L^2(\Omega)}$$

An HHO scheme

We consider the following scheme: Find $\underline{u}_h \in \underline{V}_{h,0}^k$ s.t.

$$a_h(\underline{u}_h, \underline{v}_h) := \sum_{T \in \mathcal{T}_h} a_T(\underline{u}_T, \underline{v}_T) = \int_{\Omega} f v_h \quad \forall \underline{v}_h \in \underline{V}_{h,0}^k$$

where, for all $T \in \mathcal{T}_h$,

$$a_T(\underline{u}_T, \underline{v}_T) := \int_T \nabla r_T^{k+1} \underline{u}_T \cdot \nabla r_T^{k+1} \underline{v}_T + s_T(\underline{u}_T, \underline{v}_T)$$

and the symmetric semi-definite bilinear form s_T satisfies

$$\|\underline{v}_T\|_{1,T} \lesssim a_T(\underline{v}_T, \underline{v}_T)^{\frac{1}{2}} \lesssim \|\underline{v}_T\|_{1,T} \quad \forall \underline{v}_T \in \underline{V}_T^k \tag{ST1}$$

Stability analysis

Lemma (Well-posedness of the HHO discrete problem)

The HHO problem admits a unique solution that satisfies

$$\|\underline{u}_h\|_{1,h} \lesssim \|f\|_{L^2(\Omega)}.$$

- Squaring and summing (ST1) over $T \in \mathcal{T}_h$, we have

$$\|\underline{v}_h\|_{1,h}^2 \lesssim a_h(\underline{v}_h, \underline{v}_h) \quad \forall \underline{v}_h \in \underline{V}_{h,0}^k,$$

which expresses the **coercivity of a_h**

- Using the Cauchy–Schwarz and discrete Poincaré inequalities,

$$\int_{\Omega} f v_h \leq \|f\|_{L^2(\Omega)} \|v_h\|_{L^2(\Omega)} \lesssim \|f\|_{L^2(\Omega)} \|\underline{v}_h\|_{1,h}$$

- Letting $\underline{v}_h = \underline{u}_h$ above, the conclusion follows

Error analysis I

- Let $\underline{I}_h^k : H^1(\Omega) \rightarrow \underline{V}_h^k$ be s.t.

$$\underline{I}_h^k v := ((\pi_{\mathcal{P}^{k-1}(T)} v)_{T \in \mathcal{T}_h}, (\pi_{\mathcal{P}^k(F)} v)_{F \in \mathcal{F}_h}) \quad \forall v \in H^1(\Omega)$$

- We aim at estimating the error

$$\underline{e}_h := \underline{u}_h - \underline{I}_h^k u \in \underline{V}_{h,0}^k$$

- It holds, for all $\underline{v}_h \in \underline{V}_{h,0}^k$,

$$a_h(\underline{e}_h, \underline{v}_h) = a_h(\underline{u}_h, \underline{v}_h) - a_h(\underline{I}_h^k u, \underline{v}_h) = \int_{\Omega} f v_h - a_h(\underline{I}_h^k u, \underline{v}_h) =: \mathcal{E}_h(u; \underline{v}_h)$$

- A straightforward modification of the stability proof gives

$$\|\underline{e}_h\|_{1,h} \leq \sup_{\underline{v}_h \in \underline{V}_{h,0}^k \setminus \{\underline{0}\}} \frac{\mathcal{E}_h(u; \underline{v}_h)}{\|\underline{v}_h\|_{1,h}}$$

Error analysis II

We reformulate the components of the consistency error $\mathcal{E}_h(u; \underline{v}_h)$:

$$\int_{\Omega} f v_h = - \sum_{T \in \mathcal{T}_h} \int_T \Delta u v_h = \sum_{T \in \mathcal{T}_h} \left[\int_T \nabla u \cdot \nabla v_T + \sum_{F \in \mathcal{F}_T} \int_F (\nabla u \cdot n_{TF}) (\underline{v}_F - v_T) \right]$$

$$\begin{aligned} a_h(\underline{I}_h^k u, \underline{v}_h) &= \sum_{T \in \mathcal{T}_h} \left[\int_T \nabla r_T^{k+1}(\underline{I}_T^k u) \cdot \nabla r_T^{k+1} \underline{v}_T + s_T(\underline{I}_T^k u, \underline{v}_T) \right] \\ &= \sum_{T \in \mathcal{T}_h} \left[\int_T \nabla r_T^{k+1}(\underline{I}_T^k u) \cdot \nabla v_T + \sum_{F \in \mathcal{F}_T} \int_F (\nabla r_T^{k+1}(\underline{I}_T^k u) \cdot n_{TF}) (v_F - v_T) \right] \\ &\quad + \sum_{T \in \mathcal{T}_h} s_T(\underline{I}_T^k u, \underline{v}_T) \end{aligned}$$

Error analysis III

Gathering the above results, we get

$$\begin{aligned}\mathcal{E}_h(u; \underline{v}_h) &= \sum_{T \in \mathcal{T}_h} \int_T [\nabla u - \nabla r_T^{k+1}(\underline{I}_T^k u)] \cdot \nabla v_T \\ &\quad + \sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T} \int_F [\nabla u - (\nabla r_T^{k+1}(\underline{I}_T^k u) \cdot n_{TF})] (v_F - v_T) \\ &\quad - \sum_{T \in \mathcal{T}_h} s_T(\underline{I}_T^k u, \underline{v}_T) =: \mathfrak{T}_1 + \mathfrak{T}_2 + \mathfrak{T}_3\end{aligned}$$

Approximation properties of the potential reconstruction I

- By definition, for all $T \in \mathcal{T}_h$, all $v \in H^1(T)$, and all $w \in \mathcal{P}^{k+1}(T)$,

$$\int_T \nabla r_T^{k+1}(\underline{I}_T^k v) \cdot \nabla w = - \int_T \pi_{\mathcal{P}^{k-1}(T)} v \Delta w + \sum_{F \in \mathcal{F}_T} \int_F \pi_{\mathcal{P}^k(F)} v (\nabla w \cdot n_{TF})$$

- Noticing that $\Delta w \in \mathcal{P}^{k-1}(T)$ and $\nabla w \cdot n_{TF} \in \mathcal{P}^k(F)$, we can remove the projectors and integrate by parts to obtain

$$\int_T \nabla r_T^{k+1}(\underline{I}_T^k v) \cdot \nabla w = \int_T \nabla v \cdot \nabla w \quad \forall w \in \mathcal{P}^{k+1}(T)$$

- This shows that $\nabla r_T^{k+1} \circ \underline{I}_T^k = \pi_{\nabla \mathcal{P}^{k+1}(T)} \circ \nabla$
- It can be proved that $\pi_{\nabla \mathcal{P}^{k+1}(T)} \circ \nabla$ optimally approximates gradients

Approximation properties of the potential reconstruction II

- Noticing that $\nabla v_T \in \nabla \mathcal{P}^k(T) \subset \nabla \mathcal{P}^{k+1}(T)$,

$$\int_T [\nabla u - \nabla r_T^{k+1}(\underline{I}_T^k u)] \cdot \nabla v_T = \int_T [\nabla u - \pi_{\nabla \mathcal{P}^{k+1}(T)}(\nabla u)] \cdot \nabla v_T = 0$$

we infer

$$\mathfrak{T}_1 = 0$$

- Using Cauchy–Schwarz inequalities and the definition of $\|\cdot\|_{1,h}$,

$$\mathfrak{T}_2 \leq \left(\sum_{T \in \mathcal{T}_h} h_T \|\nabla u - \pi_{\nabla \mathcal{P}^{k+1}(T)}(\nabla u)\|_{\partial T}^2 \right)^{\frac{1}{2}} \|\underline{v}_h\|_{1,h}$$

- If, additionally, $u \in H^{k+2}(\mathcal{T}_h)$,

$$\mathfrak{T}_2 \lesssim h^{k+1} |u|_{H^{k+2}(\mathcal{T}_h)} \|\underline{v}_h\|_{1,h}$$

Polynomial consistency of the stabilization I

- To have \mathfrak{T}_3 scale as \mathfrak{T}_2 , we further assume **polynomial consistency**:

$$s_T(\underline{I}_T^k w, \underline{v}_T) = 0 \quad \forall (w, \underline{v}_T) \in \mathcal{P}^{k+1}(T) \times \underline{V}_T^k \quad (\text{ST2})$$

- For all $w \in H^{k+2}(T)$, setting $|\cdot|_{s,T} := s_T(\cdot, \cdot)^{\frac{1}{2}}$, we have

$$\begin{aligned} |\underline{I}_T^k w|_{s,T} &\stackrel{(\text{ST2})}{=} \min_{v \in \mathcal{P}^{k+1}(T)} |\underline{I}_T^k(w - v)|_{s,T} \\ &\stackrel{(\text{ST1})}{\lesssim} \min_{v \in \mathcal{P}^{k+1}(T)} \|\underline{I}_T^k(w - v)\|_{1,T} \lesssim h_T^{k+1} |w|_{H^{k+2}(T)} \end{aligned}$$

hence, by Cauchy–Schwarz inequalities and again (ST1),

$$\mathfrak{T}_3 \lesssim h^{k+1} |u|_{H^{k+2}(\mathcal{T}_h)} \|\underline{v}_h\|_{1,h}$$

Polynomial consistency of the stabilization II

Theorem (Error estimate for the HHO scheme)

Denote by $u \in H_0^1(\Omega)$ the solution to the Poisson problem and by $\underline{u}_h \in \underline{V}_h^k$ its HHO approximation. Then, under (ST1)–(ST2), and further assuming $u \in H^{k+2}(\mathcal{T}_h)$, it holds

$$\|\underline{u}_h - \underline{I}_h^k u\|_{1,h} \lesssim h^{k+1} |u|_{H^{k+2}(\mathcal{T}_h)}.$$

An example of stabilization bilinear form

Example

Let, for all $T \in \mathcal{T}_h$ and all $\underline{v}_T \in \underline{V}_T^k$,

$$(\delta_T^k \underline{v}_T, (\delta_{TF}^k \underline{v}_T)_{F \in \mathcal{F}_T}) := \underline{v}_T - \underline{I}_T^k(r_T^{k+1} \underline{v}_T).$$

The stabilization bilinear form

$$s_T(\underline{w}_T, \underline{v}_T) := h_T^{-2} \int_T \delta_T^k \underline{w}_T \delta_T^k \underline{v}_T + h_T^{-1} \sum_{F \in \mathcal{F}_T} \int_F \delta_{TF}^k \underline{w}_T \delta_{TF}^k \underline{v}_T$$

satisfies properties (ST1)–(ST2).

Numerical example




Figure: $\|\underline{e}_h\|_{1,h}$ and $\|e_h\|_{L^2(\Omega)}$ as a function of h for uniformly refined triangular (top) and hexagonal (bottom) mesh families

Outline

- 1 Preliminaries
- 2 A non-conforming finite element scheme on standard meshes
- 3 An Hybrid High-Order scheme on polytopal meshes
- 4 Fully discrete analysis framework

An abstract problem and its discretization

Find $u \in U$ s.t. $a(u, v) = l(v)$ for all $v \in U$

(Π)

- U Hilbert space
- $a : U \times U \rightarrow \mathbb{R}$ continuous bilinear form
- $l : U \rightarrow \mathbb{R}$ continuous linear form

Find $u_h \in U_h$ s.t. $a_h(u_h, v_h) = l_h(v_h)$ for all $v_h \in U_h$

(Π_h)

- (finite-dimensional) vector space U_h with norm $\|\cdot\|_{U,h}$
- $a_h : U_h \times U_h \rightarrow \mathbb{R}$ bilinear form
- $l_h : U_h \rightarrow \mathbb{R}$ linear form

Approximation error

- We want to compare u and u_h without assuming that $U_h \subset U$
- The difference $u - u_h$ does not make sense in general
- Instead, we can consider

$u - R_h u_h$ with $R_h : U_h \rightarrow U$ reconstruction

or

$u_h - I_h u$ with $I_h : U \rightarrow U_h$ interpolator

- We will explore the second choice, which leads to a simpler framework

Stability and a priori estimate I

Definition (Inf-sup stability)

The bilinear form a_h is **inf-sup stable** if there is $\gamma > 0$ s.t.

$$\gamma \|w_h\|_{U,h} \leq \sup_{v_h \in U_h \setminus \{0\}} \frac{a_h(w_h, v_h)}{\|v_h\|_{U,h}} \quad \forall w_h \in U_h. \quad (\text{inf-sup})$$

Remark (Uniform inf-sup stability)

One typically requires, in practice, γ independent of h .

Stability and a priori estimate II

Lemma (A priori estimate)

If a_h is inf-sup stable, $b_h : U_h \rightarrow \mathbb{R}$ is a linear form, and $w_h \in U_h$ is s.t.

$$a_h(w_h, v_h) = b_h(v_h) \quad \forall v_h \in U_h,$$

then

$$\|w_h\|_{U,h} \leq \gamma^{-1} \|b_h\|_{U,h,*}.$$

Proof.

It suffices to write

$$\gamma \|w_h\|_{U,h} \stackrel{\text{(inf-sup)}}{\leq} \sup_{v_h \in U_h \setminus \{0\}} \frac{a_h(w_h, v_h)}{\|v_h\|_{U,h}} = \sup_{v_h \in U_h \setminus \{0\}} \frac{b_h(v_h)}{\|v_h\|_{U,h}} =: \|b_h\|_{U,h,*}. \quad \square$$

Error equation

- We want to characterize $e_h := u_h - I_h u$ as the solution to a problem
- Using linearity, we have, for all $v_h \in U_h$,

$$a_h(e_h, v_h) = a_h(u_h, v_h) - a_h(I_h u, v_h) \stackrel{(\Pi_h)}{=} l_h(v_h) - a_h(I_h u, v_h)$$

- Thus, setting

$$\mathcal{E}_h(u; v_h) := l_h(v_h) - a_h(I_h u, v_h)$$

e_h is the solution to the following **error equation**:

$$a_h(e_h, v_h) = \mathcal{E}_h(u; v_h) \quad \forall v_h \in U_h$$

Third Strang Lemma I

Lemma (Basic a priori error estimate)

Assume that a_h is inf-sup stable and let $u_h \in U_h$ solve (Π_h) . Then,

$$\|e_h\|_{U_h} \leq \gamma^{-1} \|\mathcal{E}_h(u; \cdot)\|_{U,h,*}.$$

Proof.

Apply the stability result with $w_h \leftarrow e_h$ and $b_h \leftarrow \mathcal{E}_h(u; \cdot)$. □

Definition (Consistency)

We say that **consistency** holds if

$$\|\mathcal{E}_h(u; \cdot)\|_{U,h,*} \rightarrow 0 \text{ as } h \rightarrow 0.$$

Quasi-optimality of the error estimate

- Assume that both the following quantities are **independent of h** :

$$\gamma \text{ and } \alpha := \sup_{w_h, v_h \in U_h \setminus \{0\}} \frac{a_h(w_h, v_h)}{\|w_h\|_{U,h} \|v_h\|_{U,h}}$$

- Then, we have

$$\alpha^{-1} \|\mathcal{E}_h(u; \cdot)\|_{U,h,*} \leq \|e_h\|_{U_h} \leq \gamma^{-1} \|\mathcal{E}_h(u; \cdot)\|_{U,h,*}$$

- This expresses the **quasi-optimality** of the error estimate

Key properties and their role

- Uniform inf-sup stability \implies basic error estimate
- Consistency \implies convergence
- Uniform continuity \implies quasi-optimality of the estimate



Funded by the European Union (ERC Synergy, NEMESIS, project number 101115663). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them.

Thank you for your attention!

<https://erc-nemesis.eu/events/workshop-montpellier>

References I

-  Bonaldi, F., Di Pietro, D. A., Droniou, J., and Hu, K. (2023).
An exterior calculus framework for polytopal methods.
<http://arxiv.org/abs/2303.11093>.
-  Crouzeix, M. and Raviart, P.-A. (1973).
Conforming and nonconforming finite element methods for solving the stationary Stokes equations.
RAIRO Modél. Math. Anal. Num., 7(3):33–75.
-  Di Pietro, D. A. and Droniou, J. (2018).
A third Strang lemma for schemes in fully discrete formulation.
Calcolo, 55(40).
-  Di Pietro, D. A. and Droniou, J. (2020).
The Hybrid High-Order method for polytopal meshes.
Number 19 in Modeling, Simulation and Application. Springer International Publishing.
-  Di Pietro, D. A., Droniou, J., and Rapetti, F. (2020).
Fully discrete polynomial de Rham sequences of arbitrary degree on polygons and polyhedra.
Math. Models Methods Appl. Sci., 30(9):1809–1855.
-  Di Pietro, D. A., Ern, A., and Lemaire, S. (2014).
An arbitrary-order and compact-stencil discretization of diffusion on general meshes based on local reconstruction operators.
Comput. Meth. Appl. Math., 14(4):461–472.
-  Droniou, J., Eymard, R., Gallouët, T., and Herbin, R. (2021).
Polyhedral Methods in Geosciences, chapter Non-conforming finite elements on polytopal meshes, pages 1–35.
Number 27 in SEMA-SIMAI. Springer.
-  Nédélec, J.-C. (1980).
Mixed finite elements in \mathbf{R}^3 .
Numer. Math., 35(3):315–341.

References II



Raviart, P. A. and Thomas, J. M. (1977).

A mixed finite element method for 2nd order elliptic problems.

In Galligani, I. and Magenes, E., editors, *Mathematical Aspects of the Finite Element Method*. Springer, New York.